



**Karolinska
Institutet**

Karolinska Institutet

<http://openarchive.ki.se>

This is a Peer Reviewed Manuscript version of the following article, accepted for publication in *Psychological Medicine*.

2017-07-07

The association between childhood relocations and subsequent risk of suicide attempt, psychiatric problems, and low academic achievement

Bramson, Lauren M; Rickert, Martin E; Class, Quetzal A; Sariaslan, Amir; Almqvist, Catarina; Larsson, Henrik; Lichtenstein, Paul; D'Onofrio, Brian M

Psychol Med. 2016 Apr;46(5):969-79.

<http://doi.org/10.1017/S0033291715002469>

<http://hdl.handle.net/10616/45974>

If not otherwise stated by the Publisher's Terms and conditions, the manuscript is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.



**Karolinska
Institutet**

This is the peer reviewed version of the following article
Psychol Med. 2016 Apr;46(5):969-79., which has
been published in final form at

<http://dx.doi.org/10.1017/S0033291715002469>

**The association between childhood relocations and
subsequent risk of suicide attempt, psychiatric
problems, and low academic achievement**

**Bramson, LM; Rickert, ME; Class, QA; Sariaslan, A;
Almqvist, C; Larsson, H; Lichtenstein, P; D'Onofrio, BM**

Access to the published version may require subscription.
Published with permission from: **Cambridge University
Press**

The Association between Childhood Relocations and Subsequent Risk of Suicide Attempt, Psychiatric Problems, and Low Academic Achievement

Lauren M. Bramson, BS^{1*}
lmbramso@indiana.edu

Martin E. Rickert, PhD¹
rickertm@indiana.edu

Quetzal A. Class, PhD²
qaiclass@iu.edu

Amir Sariaslan, MS³
amir.sariaslan@ki.se

Catarina Almqvist, PhD⁴
Catarina.Almqvist@ki.se

Henrik Larsson, PhD⁴
Henrik.Larsson@ki.se

Paul Lichtenstein, PhD⁴
Paul.Lichtenstein@ki.se

Brian M. D'Onofrio, PhD¹
bmdonofr@indiana.edu

¹Indiana University, Bloomington, ²University of Chicago, ³University of Oxford,
⁴Karolinska Institutet, Stockholm

This study was supported by the National Institute of Child Health and Development (B.M.D., HD061817); Indiana Clinical Translational Institute Project Development Teams (B.M.D.); the Swedish Council for Working Life and Social Research and the Swedish Research Council through the Swedish Initiative for Research on Microdata in the Social and Medical Sciences (C.A., 340-2013-5867).

Manuscript Word Count: 4,230 / 4,500

* Address for correspondence: Lauren Bramson, Psychological and Brain Sciences, Indiana University Bloomington, 1101 E. 10th St., Bloomington IN 47405
(Email: lmbramso@indiana.edu; Phone: 812-856-2588; Fax: 812-856-4544)

Abstract

Background: Given the frequency with which families change residences, the effects of childhood relocations have gained increasing research attention. Many researchers have demonstrated that childhood relocations are associated with a variety of adverse outcomes. However, drawing strong causal claims remains problematic due to uncontrolled confounding factors.

Methods: We utilized longitudinal, population-based Swedish registers to generate a nationally representative sample of offspring born 1983-1997 (n=1,510,463). Using Cox regression and logistic regression, we examined the risk for numerous adverse outcomes after childhood relocation while controlling for measured covariates. To account for unmeasured genetic and environmental confounds, we also compared differentially exposed cousins and siblings.

Results: In the cohort baseline model, each annual relocation was associated with risk for the adverse outcomes, including suicide attempt (HR=1.19; 95% CI, 1.19-1.20). However, when accounting for offspring and parental covariates (e.g., HR=1.08; 95% CI, 1.07-1.09), as well as genetic and environmental confounds shared by cousins (HR=1.07; 95% CI, 1.05-1.09) and siblings (e.g., HR=1.00; 95% CI, 0.97-1.04), the risk for suicide attempt attenuated. We found a commensurate pattern of results for severe mental illness, substance abuse, criminal convictions, and low academic achievement.

Conclusions: Previous research may have overemphasized the independent association between relocations and later adverse outcomes. The results suggest that the association between childhood relocations and suicide attempt, psychiatric problems, and low academic achievement is partially explained by genetic and environmental confounds correlated with relocations. This

study demonstrates the importance of using family-based, quasi-experimental designs to test plausible alternate hypotheses when examining causality.

Abstract Word Count: 247 / 250

Key words: Relocations; sibling-comparisons; behavior genetics; suicide; psychopathology; bipolar disorder; schizophrenia

Abbreviations:

SSRS = Statistics Sweden Regional Register

LISA = The Integrated Database for Labour Market Research

ICD = International Classification of Diseases

SAMS = Small Areas for Market Statistics

GPA = Grade point average

HR = Hazard ratio

OR = Odds ratio

Due to rates in developed countries demonstrating that people change residences more frequently in childhood than in adulthood (Jelleyman and Spencer, 2008), the ramifications of childhood relocation has gained increasing research attention. A growing body of literature suggests that children who have changed residences are at higher risk for adverse outcomes, including suicidal behavior (Qin *et al.*, 2009), psychiatric disorders (Adam, 2004, Jelleyman and Spencer, 2008), substance use (Brown *et al.*, 2012, Dewit, 1998), criminal convictions (Gasper *et al.*, 2010), and academic difficulties (Chen, 2013, Hutchings *et al.*, 2013). Several hypotheses exist as to why relocation is associated with adverse outcomes. Relocation may result in the severing of social ties and a disruption in familiar surroundings (Durkheim, 1951, Wray *et al.*, 2011), which could, for example, increase one's risk for subsequent suicidal behavior (Stack, 2000). Or, relocations may be a marker of confounding preexisting parental psychiatric disorders, parental unemployment, disorganized family life, or low family income (Dong *et al.*, 2005, Dupere *et al.*, 2008, Fomby and Cherlin, 2007, Jelleyman and Spencer, 2008, Leventhal and Brooks-Gunn, 2003, Pribesh and Downey, 1999). Despite the frequently assumed causal relation (Haynie *et al.*, 2006, Jelleyman and Spencer, 2008), drawing causal inferences between relocations and adverse outcomes remains problematic (Adam, 2004, Scanlon and Devine, 2001). Previous studies examining the effect of childhood relocation are limited primarily by: (1) small sample sizes that limit the statistical power and generalizability (Jelleyman and Spencer, 2008, Nock *et al.*, 2008a), (2) a lack of rigorous control of measured covariates (Jelleyman and Spencer, 2008), and (3) inferring a causal relation without sufficiently exploring plausible genetic and environmental confounding factors (Rutter, 2007).

Given the importance of rigorously examining plausible alternate explanations when causal claims have been made (Rutter, 2007, Shadish *et al.*, 2002), this paper aims to examine

the relation between childhood relocations (i.e., before age 12) and a variety of adverse outcomes by including more extensive measured covariates and comparing differentially exposed cousins and siblings using data collected in population-based, longitudinal, Swedish registers. Although previous studies have highlighted the vulnerability of relocations in adolescence (Paksarian *et al.*, 2014, Qin *et al.*, 2009), we focused specifically on childhood relocations due to the minimal literature examining this age range. We examined the extent to which relocations are independently associated with suicide attempt, severe mental illness, substance abuse, criminal convictions, and academic achievement. We hypothesized that the population-wide risk for adverse outcomes associated with relocations would be attenuated after accounting for numerous offspring (i.e., sex, birth parity, maternal smoking during pregnancy, mothers' cohabitation at offspring birth, parental age at childbearing) and parental covariates (i.e., parental educational attainment, country of origin, suicide attempt, suicide, severe mental illness, substance abuse, criminal history, and ranked, average family income) and controlling for unmeasured environmental and genetic confounds shared by cousins and siblings.

Methods

Data

After approval from the Institutional Review Boards at Indiana University and the Regional Ethical Review Board in Stockholm, Sweden, we generated a dataset linking information from ten longitudinal, population-based and healthcare registers using a personal identification number assigned to all Sweden residents at birth. The registers were: (1) The Medical Birth Register, which provides detailed birth information for more than 95% of pregnancies in Sweden beginning January 1, 1973 (Swedish Centre for Epidemiology, 2001), as

well as offspring antenatal and perinatal information; (2) The Multi-Generation Register, which can be used to determine biological and adoptive familial relationships for all individuals born after 1932 or have been living in Sweden since 1961 (Statistics Sweden, 2010); (3) The Statistics Sweden Regional Register (SSRS), which records an annual dichotomous change in residence beginning in 1983 (Statistics Sweden, 2013); (4) The Integrated Database for Labour Market Research (LISA), which provides longitudinal market labor information (e.g., income level, marital status, unemployment status, social welfare status, disability status, etc.) on all individuals over the age of 16 beginning in 1990 until the end of 2008, as well as an annual count of residential changes (Statistics Sweden, 2011); (5) The Cause of Death Register, which contains the date and contributing cause(s) of deaths since January 1, 1952 (Statistics Sweden, 2010); (6) The National Patient Register, which includes International Classification of Diseases Eight, Ninth, and Tenth Revisions (*ICD-8*, *ICD-9*, *ICD-10*) codes (Janssen and Kunst, 2004) and dates for hospital inpatient admissions since January 1, 1969 (Jacobsson, 2009); (7) The Migration Register, which records dates for immigrations to and emigrations from Sweden since September 11, 1901 (Statistics Sweden, 2010); (8) The National Crime Register, which records all criminal convictions (i.e., for both nonviolent and violent offenses), the date, and the sentence since January 1, 1973 for all citizens 15 years and older (Fazel and Grann, 2006); (9) The Education Register, which includes information since 1970 on the highest level of completed formal education, divided into seven ordinal groups from fewer than nine years of education to post-graduate education (D'Onofrio *et al.*, 2010); (10) The National School Register, which includes a continuous measure of academic performance aggregated across 16 subjects since 1989, and Swedish language tests since 1987 for all students graduating from 9th grade at

approximately 16 years of (The Swedish National Agency for Education, 2011). Additional information about the registers can be found elsewhere (e.g., D’Onofrio *et al.*, 2013a).

Sample

The population cohort in this study included individuals who were born in Sweden between January 1, 1983 and December 31, 1997. Using this time frame enabled us to define childhood relocations in terms of residential address changes occurring before the age of 12 and follow individuals on our main outcomes through 2009. Specifically, we wanted to ensure exposure to the risk factor (i.e., childhood relocations) occurred before any measured outcome to more accurately assess the independent association between relocation and later adverse outcomes (Kraemer *et al.*, 1997). In addition, previous research has demonstrated that the validity of suicide attempt and severe mental illness younger than age 12 is unknown (D’Onofrio *et al.*, 2013a).

Starting with 1,594,532 observations from the Medical Birth Register, we excluded individuals sequentially starting with stillbirths (n=3,608), offspring with any congenital malformation (n=63,790), and those with missing or invalid information for sex (n=65). We then excluded individuals who had no relocation data in SSRS (n=10,227), as well as one individual born after 1990 who was missing relocation data from LISA. Next, we dropped all recorded cases of suicide attempt (n=3,892) or suicide (n=2) that occurred before the age of 12. Due to the lack of documented information on predictor and outcomes variables during time spent outside of the country, we also excluded those who migrated before age 12 but subsequently returned to Sweden and had a later recorded suicide attempt (n=473) or suicide (n=9). Finally, we excluded individuals for whom the maternal age at childbearing was under 9 years (n=6), the age of death

was less than 0 ($n=1$), and the recorded date of suicide did not match the date of death ($n=1$). In total, we excluded 5.27% ($n=84,069$) of individuals born from 1983 to 1997, leaving a final cohort of 1,510,463 individuals, which included 2.38% ($n=35,940$) from multiple births. Table 1 presents a summary of offspring-specific characteristics for the study cohort.

Measures

Relocation Predictors

We operationally defined relocations using information recorded in two different registries: (1) SSRS and (2) LISA. Table 2 presents the distribution of SSRS and LISA relocations. The first (SSRS) began in 1983 and records relocations to and from homogenous, geographic clusters called Small Areas for Market Statistics (SAMS) areas. Statistics Sweden defines SAMS areas as either sub-divisions of municipalities nested within large municipalities or electoral districts nested within in small municipalities (Statistics Sweden, 2014). There are ~1,000 people in each SAMS area in Sweden with the exception of Stockholm, which averages ~2,000 people per area (Sundquist *et al.*, 2004). Note that data in SSRS are collected annually and can account only for a move between two different SAMS areas from one year to the next. Thus, we assumed that the number of SSRS relocations was an underestimate of the actual number of relocations for an individual and used data from a second register, LISA, for our sensitivity analyses. The LISA register contains a count of the total number of moves between different residential addresses within the same year starting in 1990. For each individual in the cohort, we used two different predictor variables indexing childhood relocations by summing each measure from 0 to 11 years of age. Because SSRS has a longer time span, we used the summed SSRS variables in the main analyses and the summed LISA variable for the sensitivity

analyses. An ROC analysis using relocations as measured using LISA data as the reference variable indicated that relocations estimated using SSRS data from 1990-2008 has both high sensitivity (0.82) and very high specificity (0.99).

Adverse Outcomes

The five main outcomes under study were: (1) suicide attempt, defined as the first uncertain or certain attempt requiring inpatient hospital admission at age 12 years or older (Tidemalm, 2008); (2) severe mental illness (i.e., either schizophrenia or bipolar disorder), defined as the first inpatient hospitalization at age 12 years or older (Lichtenstein *et al.*, 2009); (3) substance abuse, defined as the first inpatient hospitalization at age 12 years or older (D'Onofrio *et al.*, 2013c, Tidemalm, 2008); (4) criminal conviction, defined as the first criminal conviction at age 15 years or older (Fazel *et al.*, 2009a); and (5) low grade point average (GPA) in 9th grade, which consisted of a continuous measurement of GPA ranked into quintiles and further dichotomized, with the first quintile considered “low GPA” (Lambe *et al.*, 2006). Note that individuals who were born after December 31, 1994 were missing criminal conviction data, as these individuals were not yet 15 years old at the conclusion of the study. Likewise, individuals born after 1992 were not used in the analyses of low grades, which indexed performance at age 16. Finally, if offspring had not received a diagnosis or conviction within the study period, they contributed person-time at risk until death, emigration, or the end date of follow-up (December 31, 2009), whichever came first. Thus, the risk period began at age 12 for all outcomes except criminal convictions, which began at age 15.

Appendix A in the Online Supplementary Material lists the *ICD* codes and provides further information about the outcomes. Appendix B shows the tetrachoric correlations among

the main outcome variables, while Appendix C lists for Kaplan-Meier estimates of the survival outcomes, including death by suicide.

Covariates

Measured offspring-specific covariates, were sex, birth parity, maternal smoking during the beginning of pregnancy, maternal cohabitation status with biological father at time of offspring birth, and parental age at childbearing. The parental-specific covariates for this study were highest level of completed formal education, country of birth, death by suicide, and lifetime history of severe mental illness (i.e., either schizophrenia or bipolar disorder), suicide attempt, inpatient substance abuse, and criminal convictions. Details of all measured covariates are provided in Table 1. We additionally controlled for the average familial income, measured using percentile ranks that were computed annually to account for inflation and/or transient unmeasured disturbance. For those born before 1990 (i.e., the first year of LISA), calculation of ranked average familial income began in 1990 and ended when the individual turned 12. All offspring in the cohort had at least three years of income data available.

Analyses

We first calculated Pearson product-moment and tetrachoric correlations to quantify the bivariate association between number of relocations and the offspring and parental covariates. Next, for each outcome, we fit four regression models (described below) that sequentially adjusted for measured confounders and unmeasured familial confounders. We estimated all models using SAS 9.3 and Stata 13.1. We used Cox regression to estimate the effect of childhood relocations on the age at the first diagnosis or criminal conviction where age was measured in

units of years rounded to the nearest 0.001; to estimate the effect of childhood relocations on low GPA, we used binary logistic regression. Model 1 (i.e., cohort baseline model) estimated the association between the number of childhood relocations and each outcome while accounting for both sex and birth parity. Model 2 (i.e., adjusted model) additionally included all offspring- and parental-specific covariates. We used robust (sandwich) estimation, clustering on the maternal identifier, in both Model 1 and Model 2 to account for the non-independence of observations at the nuclear family level. Model 3 (i.e., fixed-effect, cousin comparison model) compared differentially exposed cousins to account for genetic and environment factors shared by cousins. Before fitting this model, we created a subset of the main cohort by identifying all pairs of offspring who had the same maternal grandmother identifier but different maternal identifiers. Each pair was then assigned a unique numeric identifier code. There were 1,271,618 observations (i.e., 635,809 distinct cousin pairs) from 530,809 distinct offspring in the cousin cohort subset. We obtained fixed effects estimates by: (1) using the cousin pair identifier as a strata variable during Cox and logistic model estimation, and (2) using robust standard errors to account for repeated data from individuals belonging to more than one cousin pair. Although Model 3 does not completely eliminate genetic and environmental confounding, it accounts for 6.25% or 12.5% of genetic contributions from half- and full-cousins, respectively (D'Onofrio *et al.*, 2013b). To account for additional genetic confounding, Model 4 (i.e., fixed-effect, sibling comparison model) compared differentially exposed sibling by stratifying on the maternal identifier. Note that the intraclass correlation (ICC) between siblings measured using SSRS-based relocations variable was considerably higher compared to cousins (i.e., 77% compared to 18%, respectively). The sample includes 425,833 discordant cousin pairs for relocations and 350,910 discordant sibling pairs.

After fitting the population and covariate-adjusted models for the survival outcomes, we evaluated the proportional-hazards assumption by testing for a nonzero slope in the linear regression of the scaled Schoenfeld residuals on time. If the covariate-specific or global tests indicated that the assumption did not hold, we fit additional models that included parameters representing the interaction between relocations and categorical age (i.e., ≤ 19 as the reference, 20-23, and ≥ 24). We also performed an extensive set of sensitivity analyses to address limitations inherent to our definitions of particular constructs and to our dataset.

Results

Table 3 presents the correlations estimates between the covariates and the relocation data from SSRS and LISA. While all the covariates were associated with the frequency of relocations, the magnitudes of correlation were largest for maternal ($r=-0.25$) and paternal ($r=-0.20$) age at childbearing, closely followed by familial average income ($r=-0.21$), maternal ($r=0.18$) and paternal criminality ($r=0.16$), and maternal ($r=0.14$) and paternal substance abuse ($r=0.12$). Thus, frequent relocations are associated with younger parental age at childbearing, lower socioeconomic status, and more parental substance use problems and criminality.

Table 4 and Figure 1 present the results of the Cox regression models predicting suicide attempt, severe mental illness, substance abuse, and criminal convictions from continuous SSRS relocations. For Model 1 (i.e., cohort baseline model), the observed hazard ratio (HR) for relocations predicting suicide attempt was 1.19 (95% CI, 1.12-1.18) indicating that each additional annual relocation increases the risk for attempting suicide by 19%. After adjusting for measured covariates in Model 2, the magnitude of the association was attenuated (HR=1.08; 95% CI, 1.07-1.09), but remained robust. The association was slightly weaker in Model 3, which

compared differentially exposed cousins (HR=1.07; 95% CI, 1.05-1.09). Only in Model 4, which compared differentially exposed siblings, was the association fully attenuated (HR=1.00; 95% CI, 0.97-1.04). A similar pattern of results was observed for severe mental illness, substance abuse, criminal convictions, and low GPA (i.e., the cohort-wide associations were attenuated when accounting for offspring and parental covariates, and attenuated even further when controlling for genetic and environmental factors shared by cousins and siblings). Appendix D provides the exact HRs and odds ratios (ORs) and 95% CIs that are presented in Figure 1. Appendix E lists the parameter estimate and standard error for each covariate for the main outcomes in Model 2 and 4 (i.e., the adjusted population and sibling comparison model, respectively).

We found no evidence from that the proportional-hazards assumption was violated for the models predicting suicide attempt and severe mental illness. By contrast, tests rejected the null hypothesis of zero slope for both criminal convictions and inpatient substance abuse, indicating that the log hazard ratio function is not constant over time. To address this further, we estimated the baseline and covariate-adjusted models including two additional parameters representing the interaction between relocations and categorical age. Although the interaction terms were statistically significant, the magnitudes of the differences in estimates for substance abuse and criminality were quite small (see Appendix F for details).

We used sensitivity analyses to examine several alternative explanations of our main findings and to address generalizability issues across different measures of the exposure and outcomes. First, we fit the full covariate-adjusted Model 2 to a subset of the data that excluded individuals who were missing a maternal grandmother identifier (n=153,515; see Appendix G) to determine whether or not those with missing maternal grandmothers systemically differed from

those without missing maternal grandmothers. Results were analogous to Model 2 on the full sample (see Appendix D). Second, we fit Models 1-3 predicting (1) death by suicide, (determined with certainty using *ICD* codes) to compare with previous literature regarding the effect of relocations and death by suicide specifically (Qin *et al.*, 2009), and (2) four continuous measures of academic performance (i.e., GPA, achievement Swedish, math, and English tests) to examine the validity of our derived dichotomous outcome indexing low GPA. The results (see Appendix H) of the models predicting suicide, GPA, and achievement test scores all mirrored the pattern of findings in the main analyses. Third, we re-estimated Models 1-3 on a subset of the cohort for the main outcomes (see Appendix I) and for death by suicide and academic performance (see Appendix J) using the LISA-based relocations variables to explore the extent to which findings using the more detailed measurement for relocations agreed with the analyses based on SSRS data. Although LISA spanned fewer years, we observed the same pattern of attenuation in risk association with relocations across models as in the main analyses. Fourth, we categorized the continuous, SSRS-based measurement of childhood relocations into three groups representing no moves (reference category), one to two moves (dummy coded 0/1), and three or more moves (dummy coded 0/1). We used the re-parameterized models to detect potential nonlinear associations between relocations and the outcome. Results for Models 1-3 using categorical relocations (see Appendix K) are consistent with those from the main analysis using a continuous representation of relocations as a predictor—the risks associated with more relocations were attenuated with additional statistical and methodological controls for confounding factors. Fifth, due to the fact that SAMS areas are sub-divisions of municipal districts, we analyzed an annual change in municipality to capture relocations across broadly-defined geographical areas (see Appendix L). Results were similar regardless of whether we

defined relocation as a change in SAMS area or municipal district. Finally, we analyzed two separate age ranges (i.e., 0-5, 6-11) to explore sensitive periods of childhood relocation (i.e., early and late childhood; see Appendix M) because relatively few studies examined developmental periods in relation to childhood relocations (Anderson *et al.*, 2014). Offspring who moved between the ages of 6-11 were at an increased risk compared to offspring who moved before age 6, although risk progressively attenuated across the four models, consistent with the results of the main analyses. In sum, the results of each sensitivity analysis were commensurate with those in the main analyses and followed a similar pattern of risk attenuation.

Conclusions

The current study sought to address the limitations of past literature by examining the associations between relocations specifically in childhood and later adverse outcomes using an extensive list of measured covariates and family-based, quasi-experimental designs (D'Onofrio *et al.*, 2013b). We first showed that an increase in the frequency of relocations was associated with numerous other risk factors. Those who relocated in childhood were more likely to have been born to young parents and have parents who abuse substances, have lower incomes, and engage in criminal activity, which supports previous research (Dohrenwend *et al.*, 1992, Karriker-Jaffe, 2013, US Census Bureau, 2013). We then utilized analyses to incorporate both measured covariates and family-based comparisons, including the comparison of differentially exposed cousins and siblings. Overall, the findings suggest that previous studies may have overstated the magnitude of the independent association between relocation and later adverse outcomes. Instead, our findings suggest that the associations are more likely due to: (1) the correlates of relocation (Jelleyman and Spencer, 2008) and (2) genetic and shared environmental factors

associated with suicide attempt, severe mental illness, substance abuse, criminal convictions, and academic achievement (Bondy *et al.*, 2006, Brent and Melhem, 2008, Kendler *et al.*, 2012, Lichtenstein *et al.*, 2009). The risk for adverse outcomes attenuated when controlling for genetic and shared environmental confounds, regardless of the sensitive period in which a child moved. However, future studies should further explore the role of sensitive periods when examining risk factors for a variety of adverse outcomes.

This study was strengthened by the use of multiple approaches and designs to more precisely examine the independent associations between relocation and suicide attempt, psychiatric problems, and low academic achievement. The ICCs between cousins and siblings (i.e., 18% compared to 77%, respectively) demonstrate the likelihood that siblings move together within family units, and thus siblings are less likely to be differentially exposed to relocations. Thus, the cousin comparison model strengthens the external validity of the findings (i.e., the siblings may not be representative of the entire population), while the sibling comparison model strengthens the internal validity because the design controls for more genetic and environmental factors (D'Onofrio *et al.*, 2013b). Additionally, the large sample size enabled us to estimate associations for outcomes with low base rates (Nock *et al.*, 2008a, Saha *et al.*, 2005) and provide a more diverse cross-section of the population (Adam, 2004). The inclusion of extensive sensitivity analyses reduced the possibility that our results were influenced by our construct definitions or limitations inherent to the registers. For example, results were unaffected regardless of our definition of relocations. In addition, this study examined relocations before age 12, thus expanding upon previous studies that have highlighted the vulnerability of relocations specifically in adolescence (Paksarian *et al.*, 2014, Qin *et al.*, 2009).

Conversely, there are several limitations to this study. First, the register-based measurements of the psychiatric outcomes reflect more “severe” outcomes (e.g., suicide attempt and substance abuse), although we replicated results with a variety of other outcomes that covary with suicide attempt and substance abuse (Fazel *et al.*, 2009b, Lichtenstein *et al.*, 2009, Webb *et al.*, 2011). Second, we were unable to explore moderating factors, such as the motive behind relocating (e.g., if relocation is due to a “crisis,” the effect may be more detrimental) (Tucker *et al.*, 1998), or the population density and neighborhood deprivation of the post-relocation area (Pedersen and Mortensen, 2001). Despite this, previous research has demonstrated that the association between relocations to more dense or deprived neighborhoods and schizophrenia, for example, is negligible when controlling for unmeasured familial risk factors shared by cousins and siblings (Sariaslan *et al.*, 2014). Third, the results are based solely on Swedish data, which may not generalize to other countries or cultures. However, current research suggests that the associations between risk factors and suicidal behavior, for example, are relatively consistent cross-nationally despite socioeconomic, racial, ethnic, and cultural diversity amongst a variety of countries (Nock *et al.*, 2008b). Fourth, while the main analyses demonstrate a pattern of attenuation with increased control for measured and unmeasured covariates, sibling-comparison analyses suggested that offspring who relocate more may actually be at a decreased risk for low grades (see Appendix D) but remains an open research question for future studies. However, previous literature parallels our findings suggesting that students who perform poor in school are exposed to disadvantages that better explain the relation between relocations and poor grades (Pribesh and Downey, 1999). Finally, we were unable to measure school mobility. While we analyzed moves amongst municipal districts, and thus captured a greater geographical area

possibly indicating school changes, our measure of residential mobility was only an indirect measure of school mobility.

The results suggest that clinicians should not discount relocation as a risk factor for adverse outcomes (as there is still an association in the population model) but rather refocus intervention and prevention programs. Inquiring about relocations may serve as an indicator of exposure to co-occurring risk factors (e.g., parental substance abuse). Importantly, federal programs that aid children who frequently change residences (e.g., in military families or children in foster care) focus on educational transition, such as ensuring a timely high school graduation (Anderson *et al.*, 2014). We argue that these programs should not only consider educational outcomes, but a wide range of risk factors that are associated with both relocations and other adverse outcomes.

Future studies will need to replicate this study, particularly in other countries. Additionally, because we did not include possible interactions among childhood relocations and adverse outcomes (e.g., the role of socioeconomic status), future research would benefit from these analyses. In sum, this study illustrates the importance of using rigorous research designs to test plausible alternate hypotheses, especially when studying early risk factors (D'Onofrio *et al.*, 2013b); studies that are unable to apply such designs may overestimate the magnitude of causal influences (Academy of Medical Sciences Group, 2007).

Acknowledgements

Financial Support

This study was supported by the National Institute of Child Health and Development (B.M.D., HD061817); Indiana Clinical Translational Institute Project Development Teams (B.M.D.); the Swedish Council for Working Life and Social Research and the Swedish Research Council through the Swedish Initiative for Research on Microdata in the Social and Medical Sciences (C.A., 340-2013-5867).

Conflicts of Interest

None.

Ethical Standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Academy of Medical Sciences Group** (2007). *Identifying the Environmental Causes of Disease: How Should We Decide What to Believe and When to Take Action?* Academy of Medical Sciences: London, UK.
- Adam, E. K.** (2004). Beyond Quality: Parental and Residential Stability and Children's Adjustment. *Current Directions in Psychological Science* (Wiley-Blackwell) **13**, 210-213.
- Anderson, S., Leventhal, T., Newman, S. & Dupéré, V.** (2014). Residential Mobility Among Children: A Framework for Child and Family Policy. *Cityscape* **16**, 5-36.
- Bondy, B., Buettner, A. & Zill, P.** (2006). Genetics of Suicide. *Molecular Psychiatry* **11**, 336-351.
- Brent, D. & Melhem, N.** (2008). Familial Transmission of Suicidal Behavior. *Psychiatric Clinicals of North America* **31**, 157-177.
- Brown, D., Benzeval, M., Gayle, V., Macintyre, S., O'Reilly, D. & Leyland, A. H.** (2012). Childhood Residential Mobility and Health in Late Adolescence and Adulthood: Findings from the West of Scotland Twenty-07 Study. *Journal of Epidemiology & Community Health* **66**, 942-950.
- Chen, J.** (2013). Housing Tenure, Residential Mobility and Adolescents' Education Achievement: Evidence from Sweden. *The Annals of Regional Science* **50**, 275-294.
- D'Onofrio, B. M., Class, Q. A., Rickert, M. E., Larsson, H., Långström, N. & Lichtenstein, P.** (2013a). Preterm Birth and Mortality and Morbidity: A Population-Based Quasi-Experimental Study. *JAMA Psychiatry* **70**, 1231-1240.
- D'Onofrio, B. M., Lahey, B. B., Turkheimer, E. & Lichtenstein, P.** (2013b). Critical Need for Family-Based, Quasi-Experimental Designs in Integrating Genetic and Social Science Research. *American Journal of Public Health* **103**, S46-S55.
- D'Onofrio, B. M., Rickert, M. E., Långström, N., Donahue, K. L., Coyne, C. A., Larsson, H., Ellingson, J. M., Van Hulle, C. A., Iliadou, A. N., Rathouz, P. J., Lahey, B. B. & Lichtenstein, P.** (2013c). Familial Confounding of the Association Between Maternal Smoking during Pregnancy and Offspring Substance Use and Problems: Converging Evidence Across Samples and Measures. *Archives of General Psychiatry* **69**, 1140-1150.
- D'Onofrio, B. M., Singh, A., Neiderhiser, J., Iliadou, A., Lambe, M., Hultman, C., Långström, N. & Lichtenstein, P.** (2010). A Quasi-Experimental Study of Maternal Smoking During Pregnancy and Offspring Academic Achievement. *Child Development* **81**, 80-100.
- Dewit, D. J.** (1998). Frequent Childhood Geographic Relocation: Its Impact on Drug Use Initiation and the Development of Alcohol and Other Drug-Related Problems Among Adolescents and Young Adults. *Addictive Behaviors* **23**, 623-634.
- Dohrenwend, B. P., Levav, I., Shrout, P. E., Schwartz, S., Naveh, G., Link, B. G., Skodol, A. E. & Stueve, A.** (1992). Socioeconomic Status and Psychiatric Disorders: The Causation-Selection Issue. *Science* **255**, 946-952.
- Dong, M., Anda, R., Felitti, V., Williamson, D., Dube, S., Brown, D. & Giles, W.** (2005). Childhood Residential Mobility and Multiple Health Risks During Adolescence and Adulthood: The Hidden Role of Adverse Childhood Experiences. *Archives of Pediatrics and Adolescent Medicine* **159**, 1104-1110.
- Dupere, V., Leventhal, T. & Lacourse, E.** (2008). Neighborhood Poverty and Suicidal Thoughts and Attempts in Late Adolescence *Psychological Medicine* **39**, 1295-1306.
- Durkheim, E.** (1951). *Suicide: A Study in Sociology*. Glencoe Free Press: Glencoe, IL.

- Fazel, S. & Grann, M.** (2006). The Population Impact of Severe Mental Illness on Violent Crime. *American Journal of Psychiatry* **163**, 1397-1403.
- Fazel, S., Grann, M., Carlström, E., Lichtenstein, P. & Långström, N.** (2009a). Risk Factors for Violent Crime in Schizophrenia: A National Cohort Study of 13,806 Patients. *Journal of Clinical Psychiatry* **70**, 362-369.
- Fazel, S., Långström, N., Hjern, A., Grann, M. & Lichtenstein, P.** (2009b). Schizophrenia, Substance Abuse, and Violent Crime. *JAMA* **301**, 2016-2023.
- Fomby, P. & Cherlin, A.** (2007). Family Instability and Child Well-Being. *American Sociological Review* **72**, 181-204.
- Gasper, J., DeLuca, S. & Estacion, A.** (2010). Coming and Going: Explaining the Effects of Residential and School Mobility on Adolescent Delinquency. *Social Science Research* **39**, 459-476.
- Haynie, D., South, S. & Bose, S.** (2006). Residential Mobility and Attempted Suicide among Adolescents: An Individual-Level Analysis. *The Sociological Quarterly* **47**, 693-721.
- Hutchings, H. A., Evans, A., Barnes, P., Demmler, J., Heaven, M., Hyatt, M. A., James-Ellison, M., Lyons, R. A., Maddocks, A., Paranjothy, S., Rodgers, S. E. & Dunstan, F.** (2013). Do Children Who Move Home and School Frequently Have Poorer Educational Outcomes in Their Early Years at School? An Anonymised Cohort Study. *PLoS ONE* **8**.
- Jacobsson, A.** (2009). In English - the National Patient Register *Socialstyrelsen*.
- Janssen, F. & Kunst, A.** (2004). ICD Coding Changes and Discontinuities in Trends in Cause-Specific Mortality in Six European Countries, 1950-1999. *Bulletin of the World Health Organization* **82**, 904-913.
- Jelleyman, T. & Spencer, N.** (2008). Residential Mobility in Childhood and Health Outcomes: A Systematic Review. *Journal of Epidemiology & Community Health* **62**, 584-592.
- Karriker-Jaffe, K. J.** (2013). Neighborhood Socioeconomic Status and Substance Use by U.S. Adults. *Drug and Alcohol Dependence* **133**, 212-221.
- Kendler, K. S., Sundquist, K., Ohlsson, H., Palmer, K., Maes, H., Winkleby, M. A. & Sundquist, J.** (2012). Genetic and Familial Environmental Influences on the Risk for Drug Abuse: A National Swedish Adoption Study. *Archives of General Psychiatry* **69**, 690-697.
- Kraemer, H. C., Kazdin, A. E., Offord, D. R., Kessler, R., Jensen, P. S. & Kupfer, D.** (1997). Coming to Terms With the Terms of Risk. *Archives of General Psychiatry* **54**, 337-343.
- Lambe, M., Hultman, C., Torráng, A., MacCabe, J. & Cnattingius, S.** (2006). Maternal Smoking During Pregnancy and School Performance at Age 15. *Epidemiology* **17**, 524-530.
- Leventhal, T. & Brooks-Gunn, J.** (2003). Moving to Opportunity: An Experimental Study of Neighborhood Effects on Mental Health. *American Journal of Public Health* **93**, 1576-1582.
- Lichtenstein, P., Yip, B. H., Bjork, C., Pawitan, Y., Cannon, T., Sullivan, P. F. & Hultman, C. M.** (2009). Common Genetic Influences for Schizophrenia and Bipolar Disorder: A Population-Based Study of 2 Million Nuclear Families. *Lancet* **373**, 234-239.
- Nock, M. K., Borges, G., Bromet, E., Cha, C., Kessler, R. & Lee, S.** (2008a). Suicide and Suicidal Behavior. *Epidemiologic Reviews* **30**, 133-154.
- Nock, M. K., Borges, G., Bromet, E. J., Alonso, J., Angermeyer, M., Beautrais, A., Bruffaerts, R., Chiu, W. T., de Girolamo, G., Gluzman, S., de Graaf, R., Gureje, O., Haro, J. M., Huang, Y., Karam, E., Kessler, R., Lepine, J. P., Levinson, D., Medina-Mora, M. E., Ono, Y., Posada-Villa, J. & Williams, D.** (2008b). Cross-National Prevalence and Risk Factors for Suicide Ideation, Plans and Attempts. *The British Journal of Psychiatry* **192**, 98-105.

- Paksarian, D., Eaton, W. W., Mortensen, P. B. & Pedersen, C. B.** (2014). Childhood Residential Mobility, Schizophrenia, and Bipolar Disorder: A Population-Based Study in Denmark. *Schizophrenia Bulletin*, 1-9.
- Pedersen, C. B. & Mortensen, P. B.** (2001). Evidence of a Dose-Response Relationship Between Urbanicity During Upbringing and Schizophrenia Risk. *Archives of General Psychiatry* **58**, 1039-1046.
- Pribesh, S. & Downey, D. B.** (1999). Why Are Residential and School Moves Associated with Poor School Performance? *Demography* **36**, 521-534.
- Qin, P., Mortensen, P. & Pedersen, C.** (2009). Frequent Change of Residence and Risk of Attempted and Completed Suicide Among Children and Adolescents. *Archives of General Psychiatry* **66**, 628-632.
- Rutter, M.** (2007). Proceeding From Observed Correlation to Causal Inference. *Perspectives on Psychological Science* **2**, 377-395.
- Saha, S., Chant, D., Welham, J. & McGrath, J.** (2005). A Systematic Review of the Prevalence of Schizophrenia. *PLoS ONE* **2**, e141.
- Sariaslan, A., Larsson, H., D'Onofrio, B., Långström, N., Fazel, S. & Lichtenstein, P.** (2014). Does Population Density and Neighborhood Deprivation Predict Schizophrenia? A Nationwide Swedish Quasi-Experimental Study of 2.4 Million Individuals. *Schizophrenia Bulletin* **sbu105**.
- Scanlon, E. & Devine, K.** (2001). Residential Mobility and Youth Well-Being: Research, Policy, and Practice Issues. *Journal of Sociology and Social Welfare* **28**, 119-138.
- Shadish, W. R., Cook, T. D. & Campbell, D. T.** (2002). *Experimental and Quasi-Experimental Designs for Generalized Causal Inference* / William R. Shadish, Thomas D. Cook, Donald T. Campbell. Houghton Mifflin: Boston, MA.
- Stack, S.** (2000). Suicide: A 15-Year Review of the Sociological Literature Part II: Modernization and Social Integration Perspectives. *Suicide and Life-Threatening Behavior* **30**, 163-176.
- Statistics Sweden** (2010). Multi-Generation Register 2009: A Description of Contents and Quality Orebro, Sweden.
- Statistics Sweden** (2011). Integrated Database for Labour Market Research 1990-2009.
- Statistics Sweden** (2013). Statistics Sweden Regional Register.
- Statistics Sweden** (2014). Statistics Sweden's Market Profiles: Product Catalogue 2014. (ed. S. Sweden): Stockholm.
- Sundquist, K., Winkleby, M., Ahlén, H. & Johansson, S.-E.** (2004). Neighborhood Socioeconomic Environment and Incidence of Coronary Heart Disease: A Follow-up Study of 25,319 Women and Men in Sweden. *American Journal of Epidemiology* **157**, 655-662.
- Swedish Centre for Epidemiology** (2001). The Swedish Medical Birth Register: A Summary of Content and Quality. (ed. N. B. o. H. a. Welfare): Stockholm, Sweden.
- The Swedish National Agency for Education** (2011). The Swedish National Agency for Education.
- Tidemalm, D., Långström, N., Lichtenstein, P. & Runeson, B.** (2008). Risk of Suicide After Suicide Attempt According to Coexisting Psychiatric Disorder: Swedish Cohort Study with Long Term Follow-Up. *British Medical Journal* **337**, 1328-1331.
- Tucker, C. J., Marx, J. & Long, L.** (1998). "Moving On": Residential Mobility and Children's School Lives. *Sociology of Education* **71**, 111-129.

US Census Bureau (2013). About 36 Million Americans Moved in the Last Year, Census Bureau Reports. p. 1: Washington, DC.

Webb, R., Långström, N., Runeson, B., Lichtenstein, P. & Fazel, S. (2011). Violent Offending and IQ Level as Predictors of Suicide in Schizophrenia: National Cohort Study. *Schizophrenia Research* **130**, 143-147.

Wray, M., Colen, C. & Pescosolido, B. (2011). The Sociology of Suicide. *Annual Review of Sociology* **37**, 505-528.

Table 1.

Descriptive summary of offspring and parental covariates for cohort with individuals born between January 1, 1983 and December 31, 1997.

Offspring-Specific Covariates ^a		
Variable	N (%) ^b	
Female Sex	737,135 (48.80)	
Birth Parity		
First ^c	614,136 (40.66)	
Second	544,985 (36.08)	
Third	245,676 (16.26)	
Fourth or Higher	105,666 (7.00)	
Maternal Smoking During Beginning of Pregnancy		
No ^c	1,078,039 (71.37)	
Yes	340,847 (22.57)	
Missing	91,577 (6.06)	
Mother's Cohabitation Status At Time of Offspring Birth		
Living with Father ^c	1,336,498 (94.61)	
Not Living with Father	76,197 (5.39)	
Missing	97,768 (6.47)	
Father's Age at Childbearing		
≤19	9,949 (0.66)	
20-24	165,217 (10.94)	
25-29 ^c	467,115 (30.91)	
30-34	467,478 (30.95)	
35-39	254,812 (16.87)	
40-44	94,811 (6.28)	
≥45	39,351 (2.61)	
Missing	11,730 (0.78)	
Mother's Age at Childbearing		
≤19	41,181 (2.73)	
20-24	328,462 (21.75)	
25-29 ^c	561,224 (37.15)	
30-34	396,993 (26.28)	
35-39	154,352 (10.22)	
40-44	27,278 (1.81)	
≥45	973 (0.06)	
Parental-Specific Covariates ^d		
Variable	Mother	Father ^e
Highest Level of Education		
Primary/Lower (<9 yrs) ^c	20,503 (2.32)	38,374 (4.40)
Primary/Lower (9 yrs)	83,122 (9.42)	124,496 (14.11)
Upper/Secondary (1-2 yrs)	299,725 (33.96)	328,700 (37.24)
Upper/Secondary (3 yrs)	147,991 (16.77)	116,316 (13.18)
Post-Secondary (<3 yrs)	139,541 (15.81)	118,413 (13.42)
Post-Secondary (3+ yrs)/Post-Graduate	186,615 (21.14)	142,236 (16.12)
Missing	5,077 (0.58)	13,465 (1.53)
Born in Sweden ^c	752,539 (85.27)	742,065 (84.08)
Missing	0	7,764 (0.88)
Suicide Attempt ^f	24,049 (2.72)	18,054 (2.05)
Suicide ^f	1,049 (0.12)	3,246 (0.37)
Bipolar/Schizophrenia ^f	9,177 (1.04)	6,553 (0.74)
Inpatient Substance Abuse ^f	18,663 (2.11)	36,151 (4.10)
Criminal Conviction ^f	112,174 (12.71)	373,250 (42.29)

^a Based off 1,510,463 unique offspring born 1983-1997.

^b Denotes percentages that are rounded to nearest hundredths and thus may not equal 100.

^c Indicates the reference group. ^d Based off 882,574 distinct mothers and 882,000 distinct fathers. ^e Based on fathers of the mother's first-born child.

^f Indicates a covariate with no missing variables, and thus the reference group was "No."

Table 2.

Grouped frequency distributions of the total number of childhood relocations (ages 0-11) as measured by SSRS- and LISA-based data.

Relocations	N (%) ^a
Within-Sweden Relocations (SSRS) ^b	
0 ^d	653,090 (43.24)
1-2	651,628 (43.14)
3-5	184,759 (12.23)
6-10	20,814 (1.39)
11-12	172 (0.01)
Average Annual Moves (SD)	1.11 (1.38)
Within-Sweden Relocations (LISA) ^c	
0 ^d	224,297 (14.85)
1-2	383,423 (25.38)
3-5	180,079 (11.92)
6-10	43,361 (2.87)
11-15	4,206 (0.28)
>16	708 (0.05)
Average Annual Moves (SD)	1.88 (2.06)

^a Denotes a variable in which the percentages are rounded to the nearest hundredths and thus may not equal 100.

^b Based off 1,510,463 unique individuals born 1983-1997.

^c Based off 836,074 unique individuals born 1990-1997.

^d Reference group.

Table 3.

Sample bivariate correlations between relocations (measured by SSRS and LISA) and offspring and parental covariates.

	SSRS		LISA	
	Correlation Coefficient	Sample Size	Correlation Coefficient	Sample Size
<u>Offspring-Specific Covariates</u>				
Birth Parity	-0.10	1,510,463	-0.12	836,074
Mother Not Living With Father at Time of Offspring Birth	0.14	1,412,695	0.18	766,245
Smoking During Beginning of Pregnancy	0.14	1,418,886	0.17	792,429
<u>Maternal-Specific Covariates</u>				
Born in Sweden	-0.04	1,510,463	-0.06	836,074
Criminal Conviction	0.18	1,510,463	0.18	836,074
Suicide Attempt	0.13	1,510,463	0.14	836,074
Death by Suicide	0.01	1,510,463	0.01	836,074
Severe Psychopathology	0.06	1,510,463	0.07	836,074
Inpatient Substance Abuse	0.14	1,510,463	0.15	836,074
Age at Childbearing	-0.25	1,510,463	-0.32	836,074
Highest Level of Education	-0.07	1,494,424	-0.13	828,426
<u>Paternal-Specific Covariates</u>				
Born in Sweden	-0.06	1,498,733	-0.09	829,586
Criminal Conviction	0.16	1,510,463	0.16	836,074
Suicide Attempt	0.08	1,510,463	0.09	836,074
Death by Suicide	0.02	1,510,463	0.03	836,074
Severe Psychopathology	0.03	1,510,463	0.03	836,074
Inpatient Substance Abuse	0.12	1,510,463	0.13	836,074
Age at Childbearing	-0.20	1,498,733	-0.24	829,586
Highest Level of Education	-0.03	1,491,115	-0.08	827,604
<u>Family-Specific Covariates</u>				
Rank Average Familial Income Age 0-11	-0.21	1,507,860	-0.29	836,056

Note: All correlation coefficients $p < 0.05$ for the test $H_0: \rho = 0$. Offspring sex is not included in table.

Table 4.

The association between number of childhood relocations as measured by SSRS and suicide attempt, severe mental illness, inpatient hospitalization for substance abuse, criminal conviction and low GPA.

Outcome	Model							
	1		2		3		4	
	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>
Suicide Attempt	0.18 ^a	<0.01	0.08 ^a	<0.01	0.07 ^b	0.01	0.002 ^a	0.02
Severe Mental Illness	0.19 ^a	0.01	0.11 ^a	0.01	0.07 ^b	0.03	-0.03	0.06
Inpatient Substance Abuse	0.22 ^a	<0.01	0.14 ^a	<0.01	0.08 ^b	0.01	0.02 ^a	0.02
Criminal Convictions	0.20 ^a	<0.01	0.12 ^a	<0.01	0.07 ^b	<0.01	0.00 ^a	0.01
Low GPA	0.30 ^c	<0.01	0.14 ^c	<0.01	0.08 ^d	0.01	-0.02 ^c	0.01

^a Based off cohort size of 1,510,463.

^b Based off cohort size of 1,271,618.

^c Based off cohort size of 974,008.

^d Based off cohort size of 865,983.

Note: *b* represents the estimated regression coefficients; *SE* indicates standard error.

Online Supplementary Material - Appendix Table of Contents

Appendix A. Description of the population and healthcare registers and the *International Classification of Disease (ICD)* codes used to measure offspring and parental adverse outcomes.

Appendix B. Tetrachoric correlations among the main outcomes.

Appendix C. Kaplan-Meier estimates of all the survival outcomes analyzed in the study.

Appendix D. Main analysis measuring the association between continuous relocations (SSRS) and suicide attempt, severe mental illness, substance abuse, criminal convictions and low academic achievement (HR/OR, 95% CI) for Models 1-4.

Appendix E. The parameter estimates and standard errors of the covariates included in Model 2 and 4.

Appendix F. Results for the tests of the proportional-hazards assumptions for Models 1-2 with relocation x time interaction parameter estimates.

Appendix G. Sensitivity analysis measuring the association between continuous relocations (SSRS) and suicide attempt, severe mental illness, substance abuse, criminal convictions, and low academic achievement excluding individuals with missing maternal grandmother identification numbers.

Appendix H. Sensitivity analysis measuring the association between continuous relocations (SSRS) and additional outcomes (i.e., suicide and academic achievement) for Models 1-3.

Appendix I. Sensitivity analysis measuring the association between continuous relocations (LISA) and suicide attempt, severe mental illness, substance abuse, criminal convictions, and low academic achievement for Models 1-3.

Appendix J. Sensitivity analysis measuring the association between continuous relocations (LISA) and additional outcomes (i.e., suicide and academic achievement) for Models 1-3.

Appendix K. Sensitivity analysis measuring the association between categorical relocations (SSRS) and suicide attempt, severe mental illness, substance abuse, criminal convictions, and low academic achievement for Models 1-3.

Appendix L. Sensitivity analysis measuring the association between relocations (SSRS) and suicide attempt, mental illness, substance abuse, criminal convictions, and low academic achievement and a predictor based on more broadly-defined geographic areas (municipal districts) for Models 1-4.

Appendix M. Sensitivity analysis measuring the association between relocations (SSRS) and suicide attempt, mental illness, substance abuse, criminal convictions, and low academic achievement separating childhood into sensitive age periods for Model 1-4.

Appendix A.

Description of the population and healthcare registers and the International Classification of Disease (ICD) codes used to measure offspring and parental adverse outcomes.

ICD-8, ICD-9, and ICD-10 codes were used to identify suicide attempt, severe mental illness (i.e., diagnosis of either schizophrenia or bipolar disorder), and inpatient hospitalization for substance abuse. ICD codes are not applicable for criminal convictions or academic achievement.

Outcome	Data Source	Offspring Risk Period Start and End Dates	ICD Version	ICD Code	Description
Main Analyses					
Suicide Attempt (parental and offspring)	National Patient Register (NPR)	January 1, 1995- December 31, 2009	8, 9, 10	E950-E959, E980-E989, X60-X84, Y10-Y34, Y870, Y872	Certain and uncertain attempts including violent, non-violent, and other ^a
Severe Mental Illness (parental and offspring)	NPR	January 1, 1995- December 31, 2009	8, 9, 10	295, F20; 296.1, 296.3, 296.8, 296A-296E, 296W, F30-F31	Inpatient or outpatient primary diagnoses of schizophrenia or bipolar disorder, respectively
Inpatient Substance Abuse (parental and offspring)	NPR	January 1, 1995- December 31, 2009	8, 9, 10	303, 304, 305A, 305X, F10- F16 (except x.5), F18-19 (except x.5)	Alcohol or drug use (excluding nicotine) diagnosis
Criminal Convictions (parental and offspring)	National Crime Register	January 1, 1998- December 31, 2009	N/A	N/A	All violent and nonviolent criminal convictions 15 years and older
Low GPA (offspring)	National School Register	1999-2009 ^b	N/A	N/A	The sum of standardized scales (i.e., pass, pass with distinction, and pass with honors) over 16 subjects in 9 th grade

^a All suicide attempts in the main analyses occurred after age 12 after exclusions.

^b Given that our cohort starts in 1983 and GPA assessment occurs in the 9th grade, the risk period begins approximately in 1999.

Appendix B.

Tetrachoric correlations among the main outcomes.

In order to demonstrate that the outcomes often co-occur, we estimated the pairwise tetrachoric correlations among the four (0/1) coded status variables on the survival outcomes and the binary response variable indexing low GPA. All correlations $p < 0.05$ for the test $H_0: \rho = 0$ (no correction for multiple testing). The outcomes that are most highly correlated are suicide attempt with severe mental illness and inpatient substance abuse.

Outcome	Suicide Attempt	Severe Mental Illness	Inpatient Substance Abuse	Criminal Convictions	Low GPA
Suicide Attempt	1.00	-	-	-	-
Severe Mental Illness	0.50	1.00	-	-	-
Inpatient Substance Abuse	0.56	0.41	1.00	-	-
Criminal Convictions	0.28	0.18	0.42	1.00	-
Low GPA	0.26	0.18	0.34	0.46	1.00

Appendix C.

Kaplan-Meier estimates of all the survival outcomes analyzed in the study.

This table presents Kaplan-Meier estimates at ages 15, 20, and 25 separately for the population and in the cohort stratified by the number of childhood relocations. In general, the base rates for the survival outcomes increase with age and relocations. As stated in the text, the population prevalence for low GPA in the cohort is 9.44%. Among those who did not move 4.85% had low GPA, among those who moved 1-2 times 6.23% had low GPA, and among those who moved 3 or more times 3.52% had low GPA.

Outcome	Suicide Attempt N (%) ^a	Severe Mental Illness N (%) ^a	Inpatient Substance Abuse N (%) ^a	Criminal Convictions N (%) ^a	Death by Suicide N (%) ^a
Population					
15	5,741 (0.43)	259 (0.02)	4,213 (0.33)	0 (0) ^b	46 (0.00)
20	21,283 (2.06)	2,055 (0.23)	19,017 (1.92)	104,472 (10.70)	308 (0.03)
25	27,855 (3.57)	4,191 (0.81)	24,019 (3.13)	123,911 (15.28)	510 (0.08)
0 Relocations					
15	2,080 (0.35)	84 (0.01)	1,291 (0.23)	0 (0) ^b	15 (0.00)
20	7,547 (1.66)	691 (0.17)	6,078 (1.39)	35,394 (8.26)	115 (0.03)
25	9,971 (2.89)	1,470 (0.64)	7,655 (2.23)	42,812 (12.14)	181 (0.06)
1-2 Relocations					
15	2,435 (0.42)	122 (0.02)	1,848 (0.34)	0 (0) ^b	22 (0.00)
20	9,048 (2.06)	899 (0.24)	8,487 (2.02)	45,542 (10.99)	123 (0.03)
25	11,888 (3.63)	1,801 (0.82)	10,661 (3.28)	54,055 (15.79)	213 (0.08)
>3 Relocations					
15	1,226 (0.67)	53 (0.03)	1,074 (0.62)	0 (0) ^b	9 (0.00)
20	4,688 (3.39)	465 (0.40)	4,452 (3.36)	23,536 (17.80)	70 (0.06)
25	5,996 (5.70)	920 (1.38)	5,703 (5.76)	27,044 (24.27)	116 (0.13)

^a Indicates the number and percentage of individuals who experienced the outcome in the study cohort (n=1,510,463).

^b Individuals must be age 15 and older to be included in the National Crime Register.

Note: the total number of suicide attempts is 28,288, cases of severe mental illness is 4,389, cases substance abuse is 24,344, criminal convictions is 124,870, and deaths by suicide is 516.

Appendix D.

Main analysis measuring the association between continuous relocations (SSRS) and suicide attempt, severe mental illness, substance abuse, criminal convictions and low academic achievement (HR/OR, 95% CI) for Models 1-4.

The precise results from Figure 1 are presented the form of HRs and 95% CI (except for low GPA which is in the form of ORs). Individuals are at an increased risk for adverse outcomes according to Model 1 (i.e., general population model), although all HRs progressively attenuate in Model 2 (i.e., adjusted model), Model 3 (i.e., cousin comparison model), and Model 4 (i.e., sibling comparison model).

Outcome	Model							
	1 ^a		2 ^a		3 ^b		4 ^a	
	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>
Suicide Attempt	1.19	1.19-1.20	1.08	1.07-1.09	1.07	1.05-1.09	1.00	0.97-1.04
Severe Mental Illness	1.20	1.18-1.23	1.12	1.10-1.14	1.07	1.01-1.13	0.97	0.87-1.09
Inpatient Substance Abuse	1.24	1.23-1.25	1.10	1.09-1.10	1.08	1.07-1.10	1.02	0.98-1.06
Criminal Convictions	1.22	1.22-1.23	1.09	1.08-1.09	1.07	1.06-1.08	1.00	0.98-1.02
	1 ^c		2 ^c		3 ^d		4 ^c	
	<i>OR</i>	<i>95% CI</i>	<i>OR</i>	<i>95% CI</i>	<i>OR</i>	<i>95% CI</i>	<i>OR</i>	<i>95% CI</i>
Low GPA	1.35	1.34-1.35	1.15	1.15-1.16	1.10	1.09-1.11	0.98	0.96-0.99

^a Based off cohort size of 1,510,463. ^b Based off cohort size of 1,271,618. ^c Based off cohort size of 974,008.

^d Based off cohort size of 865,983.

Appendix E.

The parameter estimates and standard errors of the covariates included in Model 2 and 4.

In order to determine which covariates are statistically significant in the final model, parameter estimates and standard errors are presented for Model 2 (i.e., adjusted population model) and Model 4 (i.e., sibling comparison model) for all main outcomes. Note that maternal- and paternal-specific covariates were not estimated for Model 4, as these variables do not differ between siblings.

	Suicide Attempt		Severe Mental Illness		Inpatient Substance Abuse	
	Model 2	Model 4	Model 2	Model 4	Model 2	Model 4
Offspring-Specific Covariates	<i>b (SE)</i>	<i>b (SE)</i>	<i>b (SE)</i>	<i>b (SE)</i>	<i>b (SE)</i>	<i>b (SE)</i>
Female Sex	0.40 (0.01)	0.45 (0.02)	0.66 (0.03)	0.64 (0.07)	0.03 (0.01)	0.10 (0.03)
Birth Order						
First ^a	-	-	-	-	-	-
Second	0.20 (0.01)	0.20 (0.03)	-0.001 (0.04)	0.18 (0.08)	0.23 (0.02)	0.36 (0.03)
Third	0.30 (0.02)	0.30 (0.05)	0.10 (0.05)	0.51 (0.15)	0.28 (0.02)	0.45 (0.06)
Fourth or Higher	0.43 (0.03)	0.48 (0.08)	0.27 (0.07)	0.82 (0.24)	0.35 (0.03)	0.63 (0.09)
Mother's Age at Childbearing						
≤19	0.19 (0.03)	-0.20 (0.10)	0.14 (0.09)	-0.44 (0.30)	0.18 (0.03)	-0.01 (0.10)
20-24	0.10 (0.02)	-0.09 (0.04)	0.04 (0.04)	-0.41 (0.15)	0.09 (0.02)	-0.003 (0.05)
25-29 ^a	-	-	-	-	-	-
30-34	-0.08 (0.02)	0.05 (0.05)	-0.01 (0.04)	0.26 (0.14)	-0.02 (0.02)	0.04 (0.05)
35-39	-0.10 (0.03)	0.23 (0.09)	0.01 (0.06)	0.37 (0.26)	-0.07 (0.03)	-0.002 (0.10)
40-44	-0.10 (0.05)	0.17 (0.19)	0.24 (0.11)	0.26 (0.50)	-0.03 (0.06)	0.09 (0.19)
≥45	0.05 (0.24)	-9.88 (132.36)	0.91 (0.38)	-	-0.04 (0.27)	0.72 (0.95)
Father's Age at Childbearing						
≤19	0.16 (0.06)	-0.17 (0.18)	-0.26 (0.17)	-0.65 (0.93)	0.19 (0.06)	0.23 (0.18)
20-24	0.08 (0.02)	-0.04 (0.05)	-0.06 (0.05)	-0.28 (0.17)	0.11 (0.02)	0.04 (0.06)
25-29 ^a	-	-	-	-	-	-
30-34	-0.02 (0.02)	0.12 (0.04)	-0.01 (0.04)	0.05 (0.13)	-0.03 (0.02)	-0.03 (0.05)
35-39	-0.01 (0.02)	0.25 (0.08)	0.11 (0.05)	0.21 (0.24)	-0.002 (0.02)	-0.08 (0.08)
40-44	0.09 (0.03)	0.39 (0.12)	0.19 (0.07)	-0.06 (0.37)	0.06 (0.03)	-0.19 (0.13)
≥45	0.14 (0.04)	0.71 (0.20)	0.40 (0.10)	0.42 (0.61)	0.23 (0.04)	-0.19 (0.21)
Missing	-0.003 (0.09)	-	-0.44 (0.25)	-	0.05 (0.09)	-
Maternal Smoking During Pregnancy	0.19 (0.01)	-0.004 (0.05)	0.12 (0.04)	-0.06 (0.15)	0.35 (0.01)	0.01 (0.05)
Missing	0.04 (0.03)	0.06 (0.06)	0.05 (0.06)	0.04 (0.16)	0.16 (0.03)	0.04 (0.06)
Mother's Cohabitation Status at Time of Offspring Birth	0.16 (0.02)	-0.01 (0.07)	0.12 (0.06)	-0.12 (0.22)	0.27 (0.02)	0.12 (0.07)
Missing	0.07 (0.03)	0.01 (0.06)	0.09 (0.07)	-0.04 (0.18)	0.13 (0.03)	0.01 (0.06)

Appendix E.

The parameter estimates and standard errors of the covariates included in Model 2 and 4 (continued).

<u>Maternal-Specific Covariates</u>						
Highest Level of Education						
Primary/Lower (<9 yrs) ^a	-	-	-	-	-	-
Primary/Lower (9 yrs)	0.20 (0.05)	-	0.30 (0.12)	-	0.23 (0.05)	-
Upper/Secondary (1-2 yrs)	0.20 (0.04)	-	0.43 (0.12)	-	0.19 (0.05)	-
Upper/Secondary (3 yrs)	0.20 (0.05)	-	0.65 (0.12)	-	0.17 (0.05)	-
Post-Secondary (<3 yrs)	0.20 (0.05)	-	0.61 (0.12)	-	0.15 (0.05)	-
Post-Secondary (3+yrs)/Post-Graduate	0.23 (0.05)	-	0.73 (0.12)	-	0.17 (0.05)	-
Missing	-0.48 (0.15)	-	0.10 (0.38)	-	0.26 (0.14)	-
Born in Sweden	0.18 (0.02)	-	0.002 (0.05)	-	-0.03 (0.02)	-
Suicide Attempt	0.46 (0.03)	-	0.46 (0.07)	-	0.34 (0.03)	-
Suicide	0.21 (0.11)	-	0.52 (0.21)	-	0.27 (0.11)	-
Severe Mental Illness	0.20 (0.04)	-	0.03 (0.08)	-	0.23 (0.04)	-
Inpatient Substance Abuse	0.19 (0.03)	-	0.03 (0.08)	-	0.33 (0.03)	-
Criminal Conviction	0.18 (0.02)	-	0.08 (0.04)	-	0.26 (0.02)	-
<u>Paternal-Specific Covariates</u>						
Highest Level of Education						
Primary/Lower (<9 yrs) ^a	-	-	-	-	-	-
Primary/Lower (9 yrs)	0.10 (0.03)	-	0.18 (0.08)	-	0.15 (0.03)	-
Upper/Secondary (1-2 yrs)	0.14 (0.03)	-	0.27 (0.08)	-	0.14 (0.03)	-
Upper/Secondary (3 yrs)	0.05 (0.03)	-	0.30 (0.08)	-	0.13 (0.04)	-
Post-Secondary (<3 yrs)	0.05 (0.03)	-	0.37 (0.08)	-	0.11 (0.04)	-
Post-Secondary (3+yrs)/Post-Graduate	-0.02 (0.03)	-	0.50 (0.08)	-	0.14 (0.04)	-
Missing	0.06 (0.07)	-	0.29 (0.17)	-	0.40 (0.07)	-
Born in Sweden	0.05 (0.02)	-	-0.09 (0.05)	-	-0.07 (0.02)	-
Suicide Attempt	0.28 (0.03)	-	0.14 (0.08)	-	0.18 (0.03)	-
Suicide	0.33 (0.06)	-	0.56 (0.14)	-	0.33 (0.07)	-
Severe Mental Illness	0.18 (0.05)	-	1.26 (0.08)	-	0.21 (0.06)	-
Inpatient Substance Abuse	0.28 (0.02)	-	0.23 (0.06)	-	0.48 (0.02)	-
Criminal Conviction	0.27 (0.01)	-	0.20 (0.03)	-	0.38 (0.01)	-
<u>Family-Specific Covariates</u>						
Average familial income 1 st quartile ^a	-	-	-	-	-	-
Average familial income 2 nd quartile	-0.04 (0.02)	-0.17 (0.07)	-0.19 (0.04)	-0.34 (0.23)	-0.07 (0.02)	-0.04 (0.07)
Average familial income 3 rd quartile	-0.24 (0.02)	-0.12 (0.09)	-0.32 (0.05)	-0.52 (0.30)	-0.13 (0.02)	0.08 (0.10)
Average familial income 4 th quartile	-3.08 (0.02)	-0.17 (0.11)	-0.50 (0.05)	-0.68 (0.36)	-0.68 (0.02)	0.05 (0.12)
Missing	-3.08 (0.71)	-	2.40 (0.77)	-	0.07 (0.72)	-

Note: Based off cohort size of 1,510,463. ^a Indicates the reference group.

Appendix E.

The parameter estimates and standard errors of the covariates included in Model 2 and 4 (continued).

Offspring-Specific Covariates	Criminal Convictions		Low Grades	
	Model 2 <i>b (SE)</i>	Model 4 <i>b (SE)</i>	Model 2 <i>b (SE)</i>	Model 4 <i>b (SE)</i>
Female Sex	-0.89 (0.01)	-0.85 (0.01)	-0.49 (0.01)	-0.64 (0.01)
Birth Order				
First ^a	-	-	-	-
Second	0.19 (0.01)	0.20 (0.01)	0.43 (0.01)	0.35 (0.02)
Third	0.26 (0.01)	0.22 (0.03)	0.75 (0.01)	0.55 (0.04)
Fourth or Higher	0.36 (0.01)	0.28 (0.04)	1.07 (0.01)	0.74 (0.05)
Mother's Age at Childbearing				
≤19	0.26 (0.02)	0.09 (0.05)	0.49 (0.02)	-0.10 (0.05)
20-24	0.13 (0.01)	0.04 (0.02)	0.23 (0.01)	-0.01 (0.03)
25-29 ^a	-	-	-	-
30-34	-0.09 (0.01)	0.01 (0.02)	-0.14 (0.01)	0.05 (0.03)
35-39	-0.15 (0.01)	0.05 (0.04)	-0.24 (0.01)	0.06 (0.06)
40-44	-0.24 (0.03)	-0.02 (0.09)	-0.30 (0.03)	0.09 (0.11)
≥45	-0.07 (0.12)	-0.14 (0.42)	-0.36 (0.13)	0.01 (0.53)
Father's Age at Childbearing				
≤19	0.14 (0.03)	0.01 (0.08)	0.35 (0.03)	0.03 (0.09)
20-24	0.09 (0.01)	0.03 (0.02)	0.17 (0.01)	-0.06 (0.03)
25-29 ^a	-	-	-	-
30-34	-0.03 (0.01)	-0.01 (0.02)	-0.05 (0.01)	-0.01 (0.03)
35-39	-0.02 (0.01)	-0.01 (0.04)	0.002 (0.01)	0.07 (0.05)
40-44	0.02 (0.02)	-0.01 (0.06)	0.10 (0.02)	0.16 (0.08)
≥45	0.06 (0.02)	0.04 (0.10)	0.18 (0.02)	0.11 (0.12)
Missing	0.03 (0.04)	-	0.14 (0.05)	-
Maternal Smoking During Pregnancy	0.28 (0.01)	-0.01 (0.02)	0.45 (0.01)	-0.03 (0.03)
Missing	0.11 (0.01)	-0.06 (0.03)	0.22 (0.01)	0.02 (0.03)
Mother's Cohabitation Status at Time of Offspring Birth	0.18 (0.01)	0.004 (0.03)	0.24 (0.01)	0.01 (0.04)
Missing	0.06 (0.01)	0.01 (0.03)	0.05 (0.01)	0.02 (0.03)

Appendix E.

The parameter estimates and standard errors of the covariates included in Model 2 and 4 (continued).

<u>Maternal-Specific Covariates</u>				
Highest Level of Education				
Primary/Lower (<9 yrs) ^a	-	-	-	-
Primary/Lower (9 yrs)	0.04 (0.02)	-	0.04 (0.02)	-
Upper/Secondary (1-2 yrs)	-0.03 (0.02)	-	-0.25 (0.02)	-
Upper/Secondary (3 yrs)	-0.08 (0.02)	-	-0.51 (0.02)	-
Post-Secondary (<3 yrs)	-0.13 (0.02)	-	-0.79 (0.02)	-
Post-Secondary (3+ yrs)/Post-Graduate	-0.15 (0.02)	-	-0.93 (0.02)	-
Missing	0.21 (0.06)	-	0.23 (0.08)	-
Born in Sweden	-0.09 (0.01)	-	-0.01 (0.01)	-
Suicide Attempt	0.14 (0.02)	-	0.20 (0.02)	-
Suicide	0.19 (0.06)	-	-0.09 (0.08)	-
Severe Mental Illness	0.04 (0.02)	-	-0.01 (0.03)	-
Inpatient Substance Abuse	0.11 (0.02)	-	0.09 (0.02)	-
Criminal Conviction	0.35 (0.01)	-	0.30 (0.01)	-
<u>Paternal-Specific Covariates</u>				
Highest Level of Education				
Primary/Lower (<9 yrs) ^a	-	-	-	-
Primary/Lower (9 yrs)	0.04 (0.01)	-	0.01 (0.02)	-
Upper/Secondary (1-2 yrs)	0.01 (0.01)	-	-0.16 (0.01)	-
Upper/Secondary (3 yrs)	-0.09 (0.02)	-	-0.48 (0.02)	-
Post-Secondary (<3 yrs)	-0.14 (0.02)	-	-0.68 (0.02)	-
Post-Secondary (3+ yrs)/Post-Graduate	-0.18 (0.02)	-	-0.78 (0.02)	-
Missing	0.08 (0.03)	-	0.01 (0.04)	-
Born in Sweden	-0.19 (0.01)	-	-0.12 (0.01)	-
Suicide Attempt	0.11 (0.02)	-	0.19 (0.02)	-
Suicide	0.16 (0.04)	-	0.13 (0.04)	-
Severe Mental Illness	0.06 (0.03)	-	-0.01 (0.04)	-
Inpatient Substance Abuse	0.22 (0.01)	-	0.31 (0.01)	-
Criminal Conviction	0.44 (0.01)	-	0.38 (0.01)	-
<u>Family-Specific Covariates</u>				
Average familial income 1 st quartile ^a	-	-	-	-
Average familial income 2 nd quartile	-0.10 (0.01)	0.02 (0.03)	-0.16 (0.01)	-0.01 (0.04)
Average familial income 3 rd quartile	-0.15 (0.01)	0.08 (0.04)	-0.32 (0.01)	-0.04 (0.05)
Average familial income 4 th quartile	-0.19 (0.01)	0.05 (0.06)	-0.52 (0.01)	-0.10 (0.06)
Missing	-1.24 (0.35)	-	-0.18 (0.25)	0.001 (1.09)

Note: Based off cohort size of 1,510,463 ^a Indicates the reference group.

Appendix F.

Results for the tests of the proportional-hazards assumptions for Models 1-2 with relocation x time interaction parameter estimates.

To test the proportionality assumption for inpatient substance abuse and criminal convictions, we included two additional parameters representing the interaction of relocations with offspring age at first diagnosis/criminal conviction. Despite the violation in the proportionality assumption, the hazard ratios differ little from models that included the relocation x time interaction and those that did not.

Outcome	Model					
	<i>b</i>	1 ^a <i>HR</i>	95% <i>CI</i>	<i>b</i>	2 ^a <i>HR</i>	95% <i>CI</i>
<u>Inpatient Substance Abuse</u>						
Without interactions	0.22*	1.24	1.23-1.25	0.09*	1.10	1.09-1.10
With interactions	0.20*	1.23	1.22-1.24	0.08*	1.08	1.07-1.09
Relocations x Age(≤19)	REF	-	-	REF	-	-
Relocations x Age(20-23)	0.05*	-	-	0.04*	-	-
Relocations x Age(≥24)	0.02*	-	-	0.02*	-	-
<u>Criminal Convictions</u>						
Without interactions	0.20*	1.22	1.22-1.23	0.08*	1.09	1.08-1.09
With interactions	0.20*	1.23	1.22-1.23	0.08*	1.09	1.08-1.09
Relocations x Age(≤19)	REF	-	-	REF	-	-
Relocations x Age(20-23)	-0.02*	-	-	-0.02*	-	-
Relocations x Age(≥24)	-0.06*	-	-	-0.06*	-	-

^a Based off cohort size of 1,510,463.

^b Based off cohort size of 1,271,618.

* Indicates significant at p<0.05.

Appendix G.

Sensitivity analysis measuring the association between continuous relocations (SSRS) and suicide attempt, severe mental illness, substance abuse, criminal convictions, and low academic achievement excluding individuals with missing maternal grandmother identification numbers.

We also estimated Model 2 (i.e., model adjusted by all individual and parental covariates) on the cohort without missing maternal grandmother identification numbers. This model yielded parameter estimates based on the same subset of the data used for the cousin comparison model. Results are similar to Model 2 in Appendix D.

Outcome	<i>HR</i> ^a	<i>95% CI</i>
Suicide Attempt	1.01	0.97-1.15
Severe Mental Illness	1.00	0.88-1.12
Inpatient Substance Abuse	1.15	1.14-1.16
Criminal Convictions	1.13	1.13-1.14
<hr/>		
	<i>OR</i> ^b	<i>95% CI</i>
Low GPA	0.98	0.96-1.00

^a Based off cohort size of 1,271,618.

^b Based off cohort size of 865,983.

Appendix H.

Sensitivity analysis measuring the association between continuous relocations (SSRS) and additional outcomes (i.e., suicide and academic achievement) for Models 1-3.

Because a large body of research has focused on relocation and suicide specifically, we performed sensitivity analyses examining suicide. To provide an additional measurement of academic achievement and further assess the main outcome of low GPA, we ran sensitivity analyses on continuous academic outcomes. The results were consistent with the finding that an individual's risk for adverse outcomes decreased when accounting for covariates and environmental and genetic confounds shared by cousins. The sibling comparison model was not included due to insufficient observations.

Outcome	Model					
	1		2		3	
	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>
Suicide	1.22 ^a	1.16-1.28	1.11 ^a	1.05-1.18	1.00 ^b	0.84-1.18
	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>
GPA	-0.14 ^c	<0.01	-0.06 ^c	<0.01	-0.05 ^d	<0.01
Achievement Swedish Test	-0.06 ^e	<0.01	-0.02 ^e	<0.01	-0.01 ^f	<0.01
Achievement Math Test	-0.08 ^g	<0.01	-0.04 ^g	<0.01	-0.02 ^h	<0.01
Achievement English Test	-0.03 ⁱ	<0.01	-0.003 ⁱ	<0.01	-0.003 ^j	<0.01

^a Based off cohort size of 1,510,463.

^c Based off cohort size of 974,008.

^e Based off cohort size of 499,116.

^g Based off cohort size of 631,908.

ⁱ Based off cohort size of 498,854.

^b Based off cohort size of 1,271,618.

^d Based off cohort size of 865,983.

^f Based off cohort size of 445,443.

^h Based off cohort size of 566,561.

^j Based off cohort size of 444,724.

Appendix I.

Sensitivity analysis measuring the association between continuous relocations (LISA) and suicide attempt, severe mental illness, substance abuse, criminal convictions, and low academic achievement for Models 1-3.

In order to validate our primary measurement of relocations (SSRS), we introduced a second measurement of relocations (LISA). Despite reduced statistical precision due to fewer relocation cases, the results lend support to the lack of an independent association from Model 1 to Model 3. The sibling comparison model was not included due to insufficient observations.

Outcome	Model					
	1 ^a		2 ^a		3 ^b	
	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>
Suicide Attempt	1.12	1.11-1.12	1.05	1.04-1.07	1.05	1.02-1.08
Severe Mental Illness	1.13	1.10-1.16	1.06	1.03-1.09	1.17	1.03-1.33
Inpatient Substance Abuse	1.15	1.14-1.15	1.09	1.08-1.10	1.09	1.05-1.12
Criminal Convictions	1.14	1.14-1.15	1.08	1.08-1.09	1.05	1.04-1.07
	1 ^c		2 ^c		3 ^d	
	<i>OR</i>	<i>95% CI</i>	<i>OR</i>	<i>95% CI</i>	<i>OR</i>	<i>95% CI</i>
Low GPA	1.26 ^a	1.25-1.26	1.11 ^a	1.10-1.11	1.08 ^b	1.06-1.10

^a Based off cohort size of 836,074.

^b Based off cohort size of 679,069.

^c Based off cohort size of 329,608.

^d Based off cohort size of 295,707.

Appendix J.

Sensitivity analysis measuring the association between continuous relocations (LISA) and additional outcomes (i.e., suicide and academic achievement) for Models 1-3.

We analyzed suicide and academic achievement using our second measurement of relocation (LISA). The risks for the outcomes decreased over the three models, paralleling previous conclusions. The sibling comparison model was not included due to insufficient observations.

Outcome	Model					
	1 ^a		2 ^a		3 ^b	
	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95%</i>
Suicide	1.12 ^a	1.05-1.19	1.07 ^a	0.99-1.05	— ^{b*}	— [*]
	1 ^c		2 ^c		3 ^d	
	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>
GPA	-0.11 ^c	<0.01	-0.11 ^c	<0.01	-0.04 ^d	0.01
Achievement Swedish Test	-0.05 ^e	<0.01	-0.02 ^e	<0.01	-0.01 ^f	<0.01
Achievement Math Test	-0.07 ^g	<0.01	-0.03 ^g	<0.01	-0.02 ^h	<0.01
Achievement English Test	-0.03 ⁱ	<0.01	-0.002 ⁱ	<0.01	0.003 ^j	<0.01

^a Based off cohort size of 836,074.

^c Based off cohort size of 329,608.

^e Based off cohort size of 409,893.

^g Based off cohort size of 400,734.

ⁱ Based off cohort size of 408,362.

^b Based off cohort size of 679,069.

^d Based off cohort size of 295,707.

^f Based off cohort size of 360,641.

^h Based off cohort size of 351,508.

^j Based off cohort size of 358,829.

* Indicates that the model could not be estimated.

Appendix K.

Sensitivity analysis measuring the association between categorical relocations (SSRS) and suicide attempt, severe mental illness, substance abuse, criminal convictions, and low academic achievement for Models 1-3.

We categorized continuous relocations measured by SSRS into three groups (i.e., 0 moves, 1-2 moves, 3 or more moves). Those who moved one or more times were at an elevated risk in Model 1 for all outcomes, although the risks attenuated in Model 2 and even further in Model 3. The sibling comparison model was not included due to insufficient observations.

Outcome	<u>Model</u>		
	1 ^a HR, 95% CI	2 ^a HR, 95% CI	3 ^b HR, 95% CI
Suicide Attempt			
0 moves	REF	REF	REF
1-2 moves	1.29 (1.25-1.32)	1.16 (1.13-1.19)	1.18 (1.12-1.25)
≥3 moves	2.10 (2.03-2.17)	1.42 (1.37-1.47)	1.39 (1.29-1.49)
Severe Mental Illness			
0 moves	REF	REF	REF
1-2 moves	1.33 (1.24-1.42)	1.20 (1.12-1.29)	1.08 (0.90-1.29)
≥3 moves	2.23 (2.06-2.42)	1.63 (1.49-1.79)	1.27 (0.99-1.61)
Inpatient Substance Abuse			
0 moves	REF	REF	REF
1-2 moves	1.52 (1.48-1.57)	1.30 (1.26-1.34)	1.18 (1.11-1.25)
≥3 moves	2.64 (2.55-2.73)	1.58 (1.52-1.64)	1.46 (1.35-1.57)
Criminal Convictions			
0 moves	REF	REF	REF
1-2 moves	1.39 (1.38-1.41)	1.20 (1.19-1.22)	1.13 (1.10-1.16)
≥3 moves	2.35 (2.31-2.38)	1.45 (1.43-1.48)	1.35 (1.30-1.40)
	1 ^c OR, 95% CI	2 ^c OR, 95% CI	3 ^d OR, 95% CI
Low GPA			
0 moves	REF	REF	REF
1-2 moves	1.54 (1.52-1.56)	1.33 (1.31-1.35)	1.25 (1.21-1.29)
≥3 moves	3.36 (3.30-3.41)	1.83 (1.80-1.87)	1.54 (1.47-1.61)

^a Based off cohort size of 1,510,463.

^b Based off cohort size of 1,271,618.

^c Based off cohort size of 974,008.

^d Based off cohort size of 865,983.

Appendix L.

Sensitivity analysis measuring the association between relocations (SSRS) and suicide attempt, mental illness, substance abuse, criminal convictions, and low academic achievement and a predictor based on more broadly-defined geographic areas (municipal districts) for Models 1-4.

When defining relocations as a change in municipality, as compared to a change in a SAMS area in the main analysis, the risk for adverse outcomes follows a similar pattern of attenuation from Model 1-4. When accounting for environmental and genetic factors shared by cousins and siblings, the risk for adverse outcomes decreases irrespective of defining relocations as a change in SAMS or municipal district.

Outcome	Model							
	1 ^a		2 ^a		3 ^b		4 ^a	
	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>	<i>HR</i>	<i>95% CI</i>
Suicide Attempt	1.21	1.19-1.22	1.08	1.07-1.10	1.07	1.04-1.10	0.96	0.91-1.01
Severe Mental Illness	1.26	1.22-1.30	1.14	1.11-1.18	1.08	1.00-1.18	1.00	0.84-1.18
Inpatient Substance Abuse	1.26	1.25-1.28	1.08	1.07-1.10	1.07	1.04-1.10	0.94	0.89-0.99
Criminal Convictions	1.23	1.23-1.24	1.08	1.07-1.08	1.07	1.06-1.09	1.00	0.98-1.03
	1 ^c		2 ^c		3 ^d		4 ^c	
	<i>OR</i>	<i>95% CI</i>	<i>OR</i>	<i>95% CI</i>	<i>OR</i>	<i>95% CI</i>	<i>OR</i>	<i>95% CI</i>
Low GPA	1.33	1.32-1.34	1.14	1.13-1.15	1.09	1.07-1.11	0.95*	0.92-0.98

^a Based off cohort size of 1,510,463.

^b Based off cohort of 1,271,618.

^c Based off cohort size of 974,008.

^d Based off cohort of 865,983.

* Indicates model did not complete Quasi-Newtonian optimization.

Appendix M.

Sensitivity analysis measuring the association between relocations (SSRS) and suicide attempt, mental illness, substance abuse, criminal convictions, and low academic achievement separating childhood into sensitive age periods for Model 1-4.

With the separation of childhood into two distinct age groups, children who relocated from age 6-11 were at an increased risk compared to relocations from birth to age 5. The risk of adverse outcomes after moving from age 6-11 followed a similar pattern of attenuation as the main analyses.

Outcome	1 ^a		2 ^a		Model 3 ^b		4 ^a	
	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI
Suicide Attempt								
0-5	1.15	1.14-1.17	1.05	1.04-1.06	1.03	1.01-1.05	0.94	0.91-0.97
6-11	1.23	1.21-1.24	1.11	1.10-1.13	1.10	1.07-1.13	0.97	0.93-1.01
Severe Mental Illness								
0-5	1.18	1.14-1.21	1.10	1.07-1.14	1.14	1.05-1.23	0.91	0.81-1.02
6-11	1.24	1.20-1.28	1.15	1.11-1.19	1.03	0.94-1.13	0.95	0.83-1.10
Inpatient Substance Abuse								
0-5	1.20	1.19-1.22	1.06	1.05-1.08	1.05	1.03-1.08	0.96	0.92-0.99
6-11	1.28	1.27-1.30	1.12	1.11-1.14	1.11	1.08-1.14	0.97	0.93-1.01
Criminal Convictions								
0-5	1.18	1.17-1.19	1.06	1.05-1.06	1.06	1.05-1.07	0.97	0.95-0.99
6-11	1.26	1.25-1.27	1.11	1.10-1.11	1.08	1.06-1.09	0.99	0.97-1.01
Low GPA	1 ^c		2 ^c		3 ^d		4 ^c	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
0-5	1.27	1.26-1.28	1.12	1.11-1.12	1.08	1.07-1.10	0.97*	0.95-0.99
6-11	1.41	1.40-1.42	1.18	1.17-1.19	1.11	1.09-1.13	0.98*	0.96-1.01

^a Based off cohort size of 1,510,463.

^d Based off cohort size of 865,983.

^b Based off cohort size of 1,271,618.

^c Based off cohort size of 974,008.

* Indicates model did not complete Quasi-Newtonian optimization.